

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
23 September 2004 (23.09.2004)

PCT

(10) International Publication Number
WO 2004/080459 A1

(51) International Patent Classification⁷: **A61K 31/445**,
A61P 25/28

(74) Agent: BUCHAN, Gavin, MacNicol; Terlings Park, East-
wick Road, Harlow, Essex CM20 2QR (GB).

(21) International Application Number:
PCT/GB2004/000983

(81) Designated States (*unless otherwise indicated, for every
kind of national protection available*): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.

(22) International Filing Date: 8 March 2004 (08.03.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/454,589 14 March 2003 (14.03.2003) US

(84) Designated States (*unless otherwise indicated, for every
kind of regional protection available*): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), Euro-
pean (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR,
GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
ML, MR, NE, SN, TD, TG).

(71) Applicants (*for all designated States except US*): MERCK
SHARP & DOHME LIMITED [GB/GB]; Hertford Road,
Hoddesdon, Hertfordshire EN11 9BU (GB). MERCK &
CO. INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ
07065 (US).

(72) Inventors; and


(75) Inventors/Applicants (*for US only*): SHEARMAN,
Mark, Steven [GB/GB]; Terlings Park, Eastwick
Road, Harlow, Essex CM20 2QR (GB). TURNER,
Mervyn [US/US]; 126 East Lincoln Avenue, Rahway, NJ
07065-0900 (US).

Published:

— with international search report

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: METHOD FOR TREATING MILD COGNITIVE IMPAIRMENT AND FOR PREVENTING OR DELAYING
ALZHEIMER'S DISEASE



(I)

(57) Abstract: There is disclosed the use of a compound of formula (I): or a pharmaceutically acceptable salt thereof, for the man-
ufacture of a medicament for the treatment of age-related cognitive decline or mild cognitive impairment, in particular with a view
to preventing or delaying the onset of Alzheimer's disease.

WO 2004/080459 A1

JC20 Rec'd PCT/PTO 1 4 SEP 2005

METHOD FOR TREATING MILD COGNITIVE IMPAIRMENT AND FOR PREVENTING OR DELAYING
ALZHEIMER'S DISEASE

This invention relates to the use of methods and materials for therapeutic treatment of the human body. In particular, it provides methods of treating age related cognitive decline and mild cognitive impairment, especially in order to prevent or
5 delay the onset of Alzheimer's disease.

Age-related cognitive decline and mild cognitive impairment (MCI) are conditions in which a memory deficit is present, but other diagnostic criteria for dementia are absent (Santacruz and Swagerty, *American Family Physician*, 63 (2001),
10 703-13). (See also "The ICD-10 Classification of Mental and Behavioural Disorders", Geneva: World Health Organisation, 1992, 64-5). When used herein, age-related cognitive decline is characterized by a decline of at least six months' duration in at least one of: memory and learning; attention and concentration; thinking; language; and visuospatial functioning and a score of more than one standard deviation below
15 the norm on standardized neuropsychologic testing such as the MMSE. In particular, there may be a progressive decline in memory. In the more severe condition mild cognitive impairment (MCI), the degree of memory impairment is outside the range considered normal for the age of the patient but Alzheimer's disease (AD) is not present. The differential diagnosis of MCI and mild AD is described by Petersen *et al.*, *Arch. Neurol.*, 56 (1999), 303-8. In the same article, Petersen *et al.* disclose that the patients suffering from MCI typically experience a progressive increase in
20 cognitive impairment and in many cases develop AD. Further information on the differential diagnosis of MCI is provided by Knopman *et al.*, *Mayo Clinic Proceedings*, 78 (2003), 1290-1308. In a study of elderly subjects, Tuokko *et al.* (*Arch. Neurol.*, 60 (2003) 577-82) found that those exhibiting MCI at the outset had a three-fold
25 increased risk of developing dementia within 5 years.

Grundman *et al.* (*J. Mol. Neurosci.*, 19 (2002), 23-28) report that lower baseline hippocampal volume in MCI patients is a prognostic indicator for subsequent AD. Similarly, Andreasen *et al.* (*Acta Neurol. Scand.*, 107 (2003) 47-51) report that high
30 CSF levels of total tau, high CSF levels of phospho-tau and lowered CSF levels of A β 42 are all associated with increased risk of progression from MCI to AD.

Age-related cognitive decline and mild cognitive impairment are distinct from the significant cognitive deficit that sometimes results from cerebral or systemic diseases and traumas, such as stroke, concussion, or major disfunction of the pituitary.

Alzheimer's disease (AD) is the most prevalent form of dementia. Its
5 diagnosis is described in the Diagnostic and Statistical Manual of Mental Disorders, 4th ed., published by the American Psychiatric Association (DSM-IV) (e.g. pages 139-143). Diagnostic criteria for Alzheimer's disease includes the development of multiple cognitive deficits in a patient manifested by both (1) memory impairment (impaired ability to learn new information or to recall previously learned information), and (2)
10 one (or more) of the following cognitive disturbances (a) aphasia (language disturbance), (b) apraxia (impaired ability to carry out motor activities despite intact motor function), (c) agnosia (failure to recognize or identify objects despite intact sensory function) and (d) disturbances in executing functioning (i.e. planning, organizing, sequencing, abstracting). Such cognitive deficits are not characterized as
15 being due to any of the following: (1) other central nervous system conditions that cause progressive deficits in memory and cognition (e.g. cerebrovascular disease, Parkinson's disease, Huntington's disease, subdural hematoma, normal-pressure hydrocephalus, brain tumor), (2) systemic conditions that are known to cause dementia (e.g. hypothyroidism, vitamin B12 or folic acid deficiency, niacin deficiency,
20 hypercalcemia, neurosyphilis, HIV infection), (3) substance-induced conditions.

Alzheimer's disease is a neurodegenerative disorder, clinically characterized by progressive loss of memory and general cognitive function, and pathologically characterized by the deposition of extracellular proteinaceous plaques in the cortical and associative brain regions of sufferers. These plaques mainly comprise fibrillar
25 aggregates of β -amyloid peptide ($A\beta$). Reducing the burden of $A\beta$ in the brain has therefore been proposed as a strategy for treatment of AD. For example Carro et al, in *Nature Medicine*, 8 (2002), 1390-7, disclose that subcutaneous administration of insulin-like growth factor 1 (IGF-1) causes a reduction in the cerebral $A\beta$ burden in certain rodents. However, some authors have questioned whether secretion of $A\beta$ is
30 responsible for the neuronal loss that is generally held to be the immediate cause of dementia of Alzheimer's type (see, for example, Robinson and Bishop, *Neurobiology of Aging*, 23 (2002), 1051-72; and also *New Scientist*, Feb. 1 2003, 35-37).

US 5,767,124, US 5,536,716, WO 94/13696 and EP 0615977B disclose compounds which are growth hormone secretagogues, i.e. which stimulate or increase the endogenous release of growth hormone in animals, including humans. This property is useful for promoting the growth of food animals, and in humans for
5 treating physiological or medical conditions characterised by a deficiency in growth hormone secretion, and medical conditions which are improved by the anabolic effects of growth hormone. The listed conditions which can be treated include Alzheimer's Disease.

US 4,902,680 advocates the administration of growth hormone to patients in
10 the advanced stages of Alzheimer's disease.


WO 00/13650 discloses that increased levels of growth hormone in the brain provide a neuroprotective effect, and in particular can rescue neurons that would otherwise die as a result of an insult such as that associated with a neurodegenerative disease. The injection of growth hormone into the brain is contemplated.

15 Major growth hormone deficiency as a result of major pituitary dysfunction can lead to cognitive impairment in adults (Deijen *et al.*, *Psychoneuroendocrinology*, 21 (1996), 313-22) which is reversible by growth hormone replacement therapy (Deijen *et al.*, *ibid.*, 23 (1998), 45-55; Soares *et al.*, *Arq. Neuropsiquiatr.*, 57 (1999), 182-9; Rosen *et al.*, *Horm. Res.*, 43 (1995), 93-99). It has also been speculated that
20 administration of growth hormone or a secretagogue thereof might, amongst many other effects, improve cognition in normal healthy elderly subjects (see, for example, Merriam *et al.*, *Endocrine*, 7 (1997), 49-52; Cummings and Merriam, *Semin. Reprod. Endocrinol.*, 17 (1999), 311-25) but no actual use was made of the secretagogues for such a purpose. However, recent publications conclude that the administration of
25 growth hormone or growth hormone secretagogues to normal ageing subjects is not of clinical benefit (Anawalt and Merriam, *Endocrinol. Metab. Clin. North Am.*, 30 (2001), 647-69; Cummings and Merriam, *Annu. Rev. Med.*, 54 (2003), 513-33). This invention is not concerned with normal healthy subjects (i.e. asymptomatic subjects).

30 A compound disclosed in the aforementioned US 5,767,124 has been the subject of a number of clinical trials in various therapeutic fields but not age-related cognitive decline or mild cognitive impairment (see, for example, Murphy *et al.*, *J. Bone Miner. Res.*, 14, (1999), 1182-8; Chapman *et al.*, *J. Clinical Endocrinology and*

Metabolism, 81, (1996), 4249-57; *ibid.*, 82, (1997), 3455-63; and Svensson et al, *ibid.*, 83, (1998), 362-9).

According to the invention, there is provided the use of a compound of formula I:



5

or a pharmaceutically acceptable salt thereof,
for the manufacture of a medicament for the treatment of age-related cognitive decline
or mild cognitive impairment. A favoured outcome of the treatment is prevention or
delay of the onset of Alzheimer's disease. Such prevention or delay may be evidenced
10 by halting or slowing of the patient's cognitive decline, or halting or slowing of the
progress of cognitive impairment.

It is one of the benefits of this invention that said medicament need not be
administrable by injection. In a preferred embodiment, said medicament is in a form
suitable for oral administration.

15 The compound of formula I may be termed N-[1(R)-[(1,2-dihydro-1-
methanesulfonylspiro[3H-indole-3,4'-piperidin]-1'-yl)carbonyl]-2-
(phenylmethoxy)ethyl]-2-amino-2-methylpropanamide.

For use in the invention, the compound of formula I is advantageously in the
form of an acid addition salt formed by interaction of the primary amine group in I
20 with a pharmaceutically acceptable acid such as hydrochloric acid, sulphuric acid,
benzenesulfonic acid, toluenesulfonic acid, methanesulfonic acid, fumaric acid, maleic
acid, succinic acid, acetic acid, benzoic acid, oxalic acid, citric acid, tartaric acid,
carbonic acid or phosphoric acid. Preferably, the compound of formula I is in the
form of the methanesulfonate salt, which is itself preferably in one of the polymorphic
25 forms described in US 5,767,124.

The synthesis of the compound of formula I and suitable salts thereof is described in US 5,767,124, EP 0615977B, US 6,028,196, US 6,046,333 and WO 96/33189.

5 The method of this invention may lead to enhanced clearance of A β from the brain, which may be evidenced by altered levels of soluble A β in the cerebrospinal fluid and/or serum. Alternatively (or additionally), imaging techniques such as magnetic resonance imaging, positron emission tomography, single photon emission computed tomography and multiphoton microscopy may be employed to monitor the extent of A β deposition in the brain (see, for example, Bacskai *et al.*, *J. Cereb. Blood*
10 *Flow Metab.*, 22 (2002), 1035-41).

In a particular embodiment, the invention provides the use of a compound as defined above for the manufacture of a medicament for preventing or delaying the onset of dementia associated with Alzheimer's disease in a patient with age related cognitive decline or in a patient with mild cognitive impairment.

15 The invention also provides a method of treating age-related cognitive decline or mild cognitive impairment comprising administering to a patient in need thereof a therapeutically-effective amount of a compound of formula I as defined above or a pharmaceutically acceptable salt thereof. In a particular embodiment, the invention also provides a method of preventing, retarding or arresting any further age-related cognitive decline
20 or progression of mild cognitive impairment comprising administering to a patient in need thereof a therapeutically-effective amount of a compound of formula I as defined above or a pharmaceutically acceptable salt thereof.

The invention also provides a method of preventing or delaying the onset of Alzheimer's disease comprising administering to a patient suffering from age related
25 cognitive decline or mild cognitive impairment a therapeutically-effective amount of a compound of formula I as defined above or a pharmaceutically acceptable salt thereof.

In a particular embodiment, the invention also provides a method of preventing or delaying the onset of dementia associated with Alzheimer's disease comprising administering to a patient suffering from age-related cognitive decline or mild
30 cognitive impairment a therapeutically-effective amount of a compound of formula I as defined above or a pharmaceutically acceptable salt thereof.

In a favoured embodiment of this invention, the compound of formula I or a pharmaceutically acceptable salt thereof is administered to a person having mild cognitive impairment.

5 In another favoured embodiment of this invention, the compound of formula I or a pharmaceutically acceptable salt thereof is administered to a person exhibiting age-related cognitive decline.

The methods of this invention are most apt for preventing, retarding or arresting the accumulation of insoluble A β in the brain of a patient suffering from age-related cognitive decline or mild cognitive impairment.

10 The medicaments useful in the invention are particularly suitable for administration to patients who suffer impaired memory function but do not exhibit other symptoms that would constitute dementia, such as aphasia, apraxia, agnosia or disturbance in executive functioning. Such impairment of memory function typically is not attributable to systemic or cerebral disease, such as stroke or metabolic disorders
15 caused by pituitary dysfunction. Such patients may be in particular people aged 55 or over, especially people aged 60 or over, and preferably people aged 65 or over. Such patients may have normal patterns and levels of growth hormone secretion for their age. However, such patients may possess one or more additional risk factors for developing Alzheimer's disease. Such factors include a family history of the disease;
20 a genetic predisposition to the disease; elevated serum cholesterol; adult-onset diabetes mellitus; raised CSF levels of total tau; raised CSF levels of phospho-tau; and lowered CSF levels of A β 42.

In a particular embodiment of the invention, the compound of formula I or a pharmaceutically acceptable salt thereof is administered to a patient suffering from
25 age-related cognitive decline or mild cognitive impairment who additionally possesses one or more risk factors for developing AD selected from: a family history of the disease; a genetic predisposition to the disease; elevated serum cholesterol; adult-onset diabetes mellitus; raised CSF levels of total tau; raised CSF levels of phospho-tau; and lowered CSF levels of A β 42.

30 A genetic predisposition (especially towards early onset Alzheimer's disease) can arise from point mutations in one or more of the APP, presenilin-1 and presenilin-

2 genes. Also, subjects who are homozygous for the $\epsilon 4$ isoform of the apolipoprotein E gene are at greater risk of developing AD.

The patient's degree of cognitive decline or impairment is advantageously assessed at regular intervals before, during and/or after a course of treatment with the compound of formula I or a pharmaceutically acceptable salt thereof, so that changes therein may be detected, e.g. the slowing or halting of cognitive decline. A variety of neuropsychological tests are known in the art for this purpose, such as the Mini-Mental State Examination (MMSE) with norms adjusted for age and education (Folstein *et al.*, *J. Psych. Res.*, 12 (1975), 196-198, Anthony *et al.*, *Psychological Med.*, 12 (1982), 397-408; Cockrell *et al.*, *Psychopharmacology*, 24 (1988), 689-692; Crum *et al.*, *J. Am. Med. Assoc'n.* 18 (1993), 2386-2391). The MMSE is a brief, quantitative measure of cognitive status in adults. It can be used to screen for cognitive decline or impairment, to estimate the severity of cognitive decline or impairment at a given point in time, to follow the course of cognitive changes in an individual over time, and to document an individual's response to treatment.

The compounds suitable for use in the invention are typically administered as pharmaceutical compositions comprising the compound of formula I (or pharmaceutically acceptable salt thereof) and a pharmaceutically acceptable carrier. Preferably these compositions are in unit dosage forms such as tablets, pills, capsules, powders, granules, sterile parenteral solutions or suspensions, metered aerosol or liquid sprays, drops, ampoules, transdermal patches, auto-injector devices or suppositories; for oral, parenteral, intranasal, sublingual or rectal administration, or for administration by inhalation or insufflation. The principal active ingredient typically is mixed with a pharmaceutical carrier, e.g. conventional tableting ingredients such as corn starch, lactose, sucrose, sorbitol, talc, stearic acid, magnesium stearate and dicalcium phosphate, or gums, dispersing agents, suspending agents or surfactants such as sorbitan monooleate and polyethylene glycol, and other pharmaceutical diluents, e.g. water, to form a homogeneous preformulation composition containing a compound of formula I or a pharmaceutically acceptable salt thereof. When referring to these preformulation compositions as homogeneous, it is meant that the active ingredient is dispersed evenly throughout the composition so that the composition may be readily subdivided into equally effective unit dosage forms such as tablets, pills and

capsules. This preformulation composition is then subdivided into unit dosage forms of the type described above, e.g. containing from 0.1 to about 150 mg of the active ingredient. Favoured unit dosage forms contain from 0.5 to 100 mg, for example 0.5, 1, 2, 3, 5, 10, 15, 20, 25, 30, 50, 60 or 75 mg, of the free base or equivalent quantities of a pharmaceutically acceptable salt thereof. Tablets or pills of the pharmaceutical composition can be coated or otherwise compounded to provide a dosage form affording the advantage of prolonged action. For example, the tablet or pill can comprise an inner dosage and an outer dosage component, the latter being in the form of an envelope over the former. The two components can be separated by an enteric layer which serves to resist disintegration in the stomach and permits the inner component to pass intact into the duodenum or to be delayed in release. A variety of materials can be used for such enteric layers or coatings, such materials including a number of polymeric acids and mixtures of polymeric acids with such materials as shellac, cetyl alcohol and cellulose acetate.

The liquid forms in which the pharmaceutical compositions may be incorporated for administration orally include aqueous solutions, liquid- or gel-filled capsules, suitably flavoured syrups, aqueous or oil suspensions, and flavoured emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil or peanut oil, as well as elixirs and similar pharmaceutical vehicles. Suitable dispersing or suspending agents for aqueous suspensions include synthetic and natural gums such as tragacanth, acacia, alginate, dextran, sodium carboxymethylcellulose, methylcellulose, poly(ethylene glycol), poly(vinylpyrrolidone) or gelatin.

Pharmaceutical formulations of the compound of formula I and suitable salts thereof are described in US 6,123,964.

For use in the invention, the optimum dosage level of the compounds defined above, in terms of safety and efficacy, may vary according to the perceived risk of developing Alzheimer's disease, and/or other factors specific to the individual patient, and may be determined by methods well known to those skilled in the art. Generally speaking, doses of about 0.01 to 5.0 mg/kg per day, preferably about 0.05 to 2.5 mg/kg per day, more preferably about 0.1 to 1.0 mg/kg of body weight per day, may be contemplated.

The compounds may be administered on any suitable regimen, for example 1, 2, 3 or 4 times per day, but administration once or twice per day is preferred, with administration once per day most preferred. Said administration is preferably oral.

The compounds may be administered at regular intervals over an extended
5 period, e.g. of 3 months, 6 months, 1 year, or more, or for the remaining lifetime of the subject.

The compound of formula I and its pharmaceutically acceptable salts are particularly suitable for oral administration. Hence, orally administrable unit dose pharmaceutical compositions are most aptly employed in the methods and uses of this
10 invention.

In one embodiment of the invention, the compound N-[1(R)-[(1,2-dihydro-1-methanesulfonylspiro[3H-indole-3,4'-piperidin]-1'-yl)carbonyl]-2-(phenylmethoxy)ethyl]-2-amino-2-methylpropanamide methanesulfonate is administered at a total dose equivalent to about 25 mg of the free base per person per
15 day. In two further embodiments said compound is administered in a total dose equivalent to 10 mg or 5 mg of free base per day. In these embodiments it is preferred that the daily dose is administered at a single time, for example as one tablet or two tablets.

Other regimens and/or dosage levels outside the limits outlined above may be
20 used if circumstances so demand.

EXAMPLES

In the following examples, the active drug is N-[1(R)-[(1,2-dihydro-1-methanesulfonylspiro[3H-indole-3,4'-piperidin]-1'-yl)carbonyl]-2-(phenylmethoxy)ethyl]-2-amino-2-methylpropanamide methanesulfonate, prepared
25 in crystalline form I as described in US 5,767,124.

Example 1 – Tablet Formulation (Wet Granulation Method)

The following formulation is used to manufacture coated tablets, each containing the
30 equivalent of 25mg free base:

<u>Tablet Core</u>	<u>mg/tablet</u>
active drug	29.6
Starch Pregelatinized NF	113.0
Calcium Phosphate Dibasic, USP	174.0
Microcrystalline Cellulose NF	57.0
Magnesium Stearate Impalpable Powder NF	2.0
Croscarmellose Sodium NF	24.0
Ethanol 95% USP*	-----
Water Purified USP*	-----

<u>Tablet Coat</u>	
Hydroxypropyl Methylcellulose USP	3.2
Hydroxypropyl Cellulose w/ 0.3% Silica	3.2
Titanium Dioxide USP	1.28
Talc USP Purified	0.32
Water Purified USP*	----

* removed during process.

The active drug, calcium phosphate, starch pregelatinized, microcrystalline cellulose and half of the croscarmellose sodium are mixed and then transferred to a granulator for brief additional mixing. A 25% ethanol/water mixture is added slowly and
5 granulation performed. The granules are dried, screened, and mixed with the remaining croscarmellose sodium and then with the magnesium stearate. Tablets are formed by compression and film coated by spraying with an aqueous suspension of the tablet coat ingredients.

Tablets containing different loadings of the drug, for example 10 mg or 5 mg,
10 are formed by the same procedure, with appropriate adjustment of the relative proportions of the ingredients.

Example 2 – Tablet formulation (Roller Compaction Process)

The following formulation is used to manufacture coated tablets, each containing the
15 equivalent of 25mg free base:

<u>Tablet Core</u>	<u>mg/tablet</u>
active drug	29.6
Calcium Phosphate Dibasic, USP	60.44
Microcrystalline Cellulose NF	150.0
Magnesium Stearate Impalpable Powder NF	2.56
Croscarmellose Sodium NF	7.5

<u>Tablet Coat</u>	
Hydroxypropyl Methylcellulose USP	3.125
Hydroxypropyl Cellulose w/ 0.3% Silica	3.125
Titanium Dioxide USP	1.25
Water Purified USP*	----

* removed during process.

The active drug, microcrystalline cellulose and a portion of the magnesium stearate are mixed, then roller compacted, and the resulting compact is milled. The milled material is blended with the croscarmellose sodium, calcium phosphate and the remaining magnesium stearate, then compressed into tablets. The tablets are film coated by spraying with an aqueous suspension of the tablet coat ingredients.

Tablets containing different loadings of the drug are formed by the same procedure, with appropriate adjustment of the relative proportions of the ingredients.

10 Example 3 – Treatment for Preventing or Delaying the Onset of Alzheimer's Disease.

One 25mg tablet (as described in Example 1 or Example 2) is administered daily with water to subjects in need of such treatment.

15 Example 4- Treatment for Preventing or Delaying the Onset of Alzheimer's Disease in a Subject Exhibiting Mild Cognitive Impairment


A subject having mild cognitive impairment is identified using the MMSE or similar diagnostic tool.

One 25mg tablet (as described in Example 1 or Example 2) is administered daily with water to said subject.

The cognitive status of the subject is monitored periodically using the MMSE or similar tool, and the subject is monitored for clinical symptoms of dementia.

CLAIMS:

1. The use of a compound of formula I:



or a pharmaceutically acceptable salt thereof for the manufacture of a medicament for treating age-related cognitive decline or mild cognitive impairment in a patient in need thereof.

2. Use of a compound of formula I as defined in claim 1 or a pharmaceutically acceptable salt thereof for the manufacture of a medicament for preventing or delaying the onset of Alzheimer's disease in a patient suffering from age-related cognitive decline or mild cognitive impairment.

3. Use of a compound of formula I as defined in claim 1 or a pharmaceutically acceptable salt thereof for the manufacture of a medicament for preventing or delaying the onset of dementia associated with Alzheimer's disease in a patient suffering from age-related cognitive decline or mild cognitive impairment.

4. Use according to any previous claim wherein said patient is a person aged 55 or over.

5. Use according to any previous claim wherein said patient is a person aged 60 or over.

6. Use according to any previous claim wherein said patient is a person aged 65 or over.

5 7. Use according to any previous claim wherein said patient additionally possesses one or more risk factors for developing Alzheimer's disease selected from: a family history of the disease; a genetic predisposition to the disease; elevated serum cholesterol; adult-onset diabetes mellitus; raised CSF levels of total tau; raised CSF levels of phospho-tau; and lowered CSF levels of A β 42.

10 8. Use according to any of claims 1-7 wherein the patient suffers from mild cognitive impairment.

9. Use according to any of claims 1-7 wherein the patient suffers from age-related cognitive decline.

15

10. Use of a compound of formula I as defined in claim 1 or a pharmaceutically acceptable salt thereof for the manufacture of a medicament for preventing, retarding or arresting the accumulation of insoluble A β in the brain of a patient suffering from age-related cognitive decline or mild cognitive impairment.

20

11. Use according to any previous claim wherein the medicament is adapted for oral administration.

INTERNATIONAL SEARCH REPORT

GB2004/000983

A. CLASSIFICATION OF SUBJECT MATTER
 IPC 7 A61K31/445 A61P25/28

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, WPI Data, EMBASE, BIOSIS, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 767 124 A (KAUFMAN MICHAEL J ET AL) 16 June 1998 (1998-06-16) cited in the application abstract column 1, last paragraph - column 2, line 14 column 2, line 48 - column 3, line 5 column 16, line 20 - column 17, line 34 column 18, line 10 - line 15	1-11
Y	-----	1-11
X	EP 0 615 977 A (MERCK & CO INC) 21 September 1994 (1994-09-21) cited in the application abstract page 10, line 1 - line 17 page 30, line 45 - line 47 page 30, line 58 ----- -/-	1-11

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

15 June 2004

Date of mailing of the international search report

28/06/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Langer, O

INTERNATIONAL SEARCH REPORT

GB2004/000983

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 1 258 250 A (KAKEN PHARMA CO LTD) 20 November 2002 (2002-11-20) paragraph '0004! paragraph '0008! paragraph '0013! paragraph '0028! - paragraph '0029! paragraph '0087! claims 1-7	1
X	WO 97/41879 A (MERCK & CO INC ; WALDSTREICHER JOANNE (US)) 13 November 1997 (1997-11-13) abstract page 1, line 26 - line 29 page 2, line 9 - line 11 page 2, line 15 - line 17 page 21, line 11 - line 16 page 26, line 7 - line 24	1
Y	SRAMEK J J ET AL: "THE STATUS OF ONGOING TRIALS FOR MILD COGNITIVE IMPAIRMENT" EXPERT OPINION ON INVESTIGATIONAL DRUGS, ASHLEY PUBLICATIONS LTD., LONDON, GB, vol. 10, no. 4, 2001, pages 741-752, XP008012341 ISSN: 1354-3784 abstract page 741, paragraph 1 page 742, left-hand column, paragraph 2 - right-hand column, paragraph 1 page 743, left-hand column, paragraph 2 - right-hand column, last paragraph page 744, left-hand column, paragraph 1 page 744, right-hand column, paragraph 1	1-11
Y	CHAPMAN, IAN M. ET AL: "Stimulation of the growth hormone (GH)-insulin-like growth factor I axis by daily oral administration of a GH secretagogue (MK-677) in healthy elderly subjects" JOURNAL OF CLINICAL ENDOCRINOLOGY AND METABOLISM , 81(12), 4249-4257 CODEN: JCEMAZ; ISSN: 0021-972X, 1996, XP002284306 abstract page 4250, left-hand column, paragraph 3 - paragraph 4 page 4256, left-hand column, paragraph 2	4-6

-/--

INTERNATIONAL SEARCH REPORT

GB2004/000983

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	SMITH R G: "The aging process: where are the drug opportunities?" CURRENT OPINION IN CHEMICAL BIOLOGY. AUG 2000, vol. 4, no. 4, August 2000 (2000-08), pages 371-376, XP002284307 ISSN: 1367-5931 page 371 - page 373	4-6
Y	PATCHETT, A. A. ET AL: "Design and biological activities of L-163,191 (MK-0677): a potent, orally active growth hormone secretagogue" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA , 92(15), 7001-5 CODEN: PNASA6; ISSN: 0027-8424, 1995, XP002284304 page 7001, right-hand column, paragraph 3 page 7005, right-hand column, paragraph 2 figure 1	11
Y	PONG, SHENG-SHUNG ET AL: "Identification of a new G-protein-linked receptor for growth hormone secretagogues" MOLECULAR ENDOCRINOLOGY , 10(1), 57-61 CODEN: MOENEN; ISSN: 0888-8809, 1996, XP002284305 abstract page 57, line R figure 1	11
A	BURNS A ET AL: "Mild cognitive impairment in older people" LANCET, XX, XX, vol. 360, no. 9349, 14 December 2002 (2002-12-14), pages 1963-1965, XP004398792 ISSN: 0140-6736 page 1964, left-hand column, paragraph 3 - paragraph 4 page 1964, right-hand column, paragraph 2 - page 1965, left-hand column, paragraph 3	1-11
A	US 4 902 680 A (AROONSAKUL CHAOVANE) 20 February 1990 (1990-02-20) cited in the application abstract column 2, line 57 - line 64 column 3, line 1 - line 28 claims 1,4	1-11

-/--

INTERNATIONAL SEARCH REPORT

GB2004/000983

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>WO 00/13650 A (CLARK ROSS GRAHAM ; WILLIAMS CHRIS EDWARD (NZ); NEURONZ LIMITED (NZ);) 16 March 2000 (2000-03-16) cited in the application abstract page 2, line 17 - line 20 page 3, line 10 - line 18 page 4, line 17 - line 20 page 5, line 1 - line 3 claims 17,19,20</p>	1-11
A	<p>US 6 028 196 A (HOUGHTON PETER ET AL) 22 February 2000 (2000-02-22) cited in the application abstract</p>	1-11
A	<p>US 6 046 333 A (MOLINA AUDREY ET AL) 4 April 2000 (2000-04-04) cited in the application abstract</p>	1-11
A	<p>WO 96/33189 A (MERCK & CO INC ; ASKIN DAVID (US); MALIGRES PETER E (US); REIDER PAUL) 24 October 1996 (1996-10-24) cited in the application abstract</p>	1-11
P,Y	<p>ANDREASEN NIELS ET AL: "Cerebrospinal fluid levels of total-tau, phospho-tau and A beta 42 predicts development of Alzheimer's disease in patients with mild cognitive impairment." ACTA NEUROLOGICA SCANDINAVICA. SUPPLEMENTUM. 2003, vol. 179, 2003, pages 47-51, XP002284308 ISSN: 0065-1427 cited in the application abstract page 47 - page 48, left-hand column, paragraph 1 page 50, left-hand column, paragraph 2</p>	7
P,A	<p>ARNÁIZ EVA ET AL: "Neuropsychological features of mild cognitive impairment and preclinical Alzheimer's disease." ACTA NEUROLOGICA SCANDINAVICA. SUPPLEMENTUM. 2003, vol. 179, 2003, pages 34-41, XP002284309 ISSN: 0065-1427 abstract page 34 page 39, right-hand column, paragraph 2</p>	1-11
	-/--	

INTERNATIONAL SEARCH REPORT

GB2004/000983

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,A	SMALL BRENT J ET AL: "Cognitive deficits in preclinical Alzheimer's disease." ACTA NEUROLOGICA SCANDINAVICA. SUPPLEMENTUM. 2003, vol. 179, 2003, pages 29-33, XP002284310 ISSN: 0065-1427 abstract page 29, left-hand column, paragraph 1 page 30, left-hand column, paragraph 2 - page 31, right-hand column, paragraph 2 -----	1-11
P,A	PALMER KATIE ET AL: "What is mild cognitive impairment? Variations in definitions and evolution of nondemented persons with cognitive impairment." ACTA NEUROLOGICA SCANDINAVICA. SUPPLEMENTUM. 2003, vol. 179, 2003, pages 14-20, XP002284311 ISSN: 0065-1427 page 14, left-hand column - right-hand column, paragraph 1 page 16, right-hand column, paragraph 1 page 17, right-hand column, paragraph 2 -----	1-11
P,A	WAHLUND LARS-OLOF ET AL: "Mild cognitive impairment: experience from a memory clinic." ACTA NEUROLOGICA SCANDINAVICA. SUPPLEMENTUM. 2003, vol. 179, 2003, pages 21-24, XP002284312 ISSN: 0065-1427 abstract -----	1-11
P,A	JELIC VESNA ET AL: "Mild cognitive impairment: unanswered questions and future directions." ACTA NEUROLOGICA SCANDINAVICA. SUPPLEMENTUM. 2003, vol. 179, 2003, pages 100-101, XP002284313 ISSN: 0065-1427 page 100, left-hand column, paragraph 1 - paragraph 2 page 101, right-hand column, paragraph 1 -----	1-11
A	EP 1 149 583 A (PFIZER PROD INC) 31 October 2001 (2001-10-31) paragraph '0001! paragraph '0003! paragraph '0039! -----	1-11

INTERNATIONAL SEARCH REPORT

GB2004/000983

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5767124	A	16-06-1998	AU 707946 B2 22-07-1999
			AU 7468696 A 15-05-1997
			BG 102476 A 30-04-1999
			BR 9611229 A 25-05-1999
			CA 2235371 A1 01-05-1997
			CN 1205703 A ,B 20-01-1999
			CZ 9801280 A3 16-09-1998
			EA 528 B1 28-10-1999
			EE 9800147 A 15-12-1998
			EP 1019402 A1 19-07-2000
			HK 1017894 A1 28-09-2001
			HU 9902208 A2 28-10-1999
			JP 10512295 T 24-11-1998
			JP 3204266 B2 04-09-2001
			NO 981867 A 29-06-1998
			NZ 321370 A 29-11-1999
			PL 327511 A1 21-12-1998
			SK 51398 A3 02-12-1998
			TR 9800726 T2 21-08-1998
			WO 9715574 A1 01-05-1997
			ZA 9608989 A 29-04-1997
EP 0615977	A	21-09-1994	US 5536716 A 16-07-1996
			AT 220071 T 15-07-2002
			AU 673552 B2 14-11-1996
			AU 5232093 A 23-06-1994
			BG 61979 B1 30-11-1998
			BG 99710 A 31-01-1996
			CA 2110670 A1 12-06-1994
			CN 1092071 A ,B 14-09-1994
			CZ 9501514 A3 13-12-1995
			DE 69332075 D1 08-08-2002
			DE 69332075 T2 16-01-2003
			DK 615977 T3 22-07-2002
			EP 0615977 A1 21-09-1994
			ES 2177538 T3 16-12-2002
			FI 952863 A 09-06-1995
			HR 931486 B1 31-08-2003
			HU 9500324 A3 28-09-1995
			HU 72076 A2 28-03-1996
			JP 2509530 B2 19-06-1996
			JP 6263737 A 20-09-1994
			KR 203962 B1 15-06-1999
			MX 9307850 A1 30-06-1994
			NO 952295 A 10-08-1995
			NZ 258412 A 29-01-1997
			PL 309331 A1 02-10-1995
			PT 615977 T 31-10-2002
			RO 115804 B1 30-06-2000
			RU 2168512 C2 10-06-2001
			SI 9300646 A 30-09-1994
			SK 75995 A3 08-11-1995
			WO 9413696 A1 23-06-1994
			US 5652235 A 29-07-1997
			AT 178070 T 15-04-1999
			AU 673017 B2 24-10-1996
			AU 5232193 A 23-06-1994
			BR 1100850 A3 07-12-1999

INTERNATIONAL SEARCH REPORT

GB2004/000983

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0615977 A		CA 2110672 A1	12-06-1994
		CN 1092767 A ,B	28-09-1994
		CZ 9501515 A3	13-12-1995
		DE 69324148 D1	29-04-1999
		DE 69324148 T2	28-10-1999
		EP 0662481 A1	12-07-1995
		ES 2129072 T3	01-06-1999
		FI 952862 A	09-06-1995
		HR 931484 A1	30-04-1996
		HU 73228 A2	29-07-1996
		IL 107836 A	04-01-1998
		JP 2509147 B2	19-06-1996
		JP 7097367 A	11-04-1995
		MX 9307851 A1	30-06-1994
EP 1258250 A	20-11-2002	AU 2405401 A	09-07-2001
		CA 2395996 A1	05-07-2001
		EP 1258250 A1	20-11-2002
		WO 0147558 A1	05-07-2001
		US 2003100494 A1	29-05-2003
WO 9741879 A	13-11-1997	AU 711884 B2	21-10-1999
		AU 2996397 A	26-11-1997
		CA 2253390 A1	13-11-1997
		EP 0900086 A1	10-03-1999
		JP 2000511885 T	12-09-2000
		WO 9741879 A1	13-11-1997
US 4902680 A	20-02-1990	CA 1323294 C	19-10-1993
		US 4898856 A	06-02-1990
		US 4791099 A	13-12-1988
		US 4727041 A	23-02-1988
		AU 1012588 A	13-07-1989
		US 5017470 A	21-05-1991
		DK 69189 A	17-08-1989
		US 4898857 A	06-02-1990
		US 4897389 A	30-01-1990
WO 0013650 A	16-03-2000	AU 757907 B2	13-03-2003
		AU 5885999 A	27-03-2000
		BR 9913457 A	16-10-2001
		CA 2343214 A1	16-03-2000
		CN 1316908 T	10-10-2001
		EP 1109572 A2	27-06-2001
		JP 2002524402 T	06-08-2002
		WO 0013650 A2	16-03-2000
		NZ 510253 A	28-02-2003
US 6028196 A	22-02-2000	AU 7468596 A	15-05-1997
		BR 9610887 A	13-07-1999
		CA 2235511 A1	01-05-1997
		EP 0863900 A1	16-09-1998
		JP 11513984 T	30-11-1999
		PL 326428 A1	14-09-1998
		SK 51198 A3	04-11-1998
		CZ 9801282 A3	15-07-1998
		HR 960494 A1	30-04-1998
		HU 9902095 A2	28-10-1999

INTERNATIONAL SEARCH REPORT

GB2004/000983

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 6028196	A		WO 9715573 A1	01-05-1997
			US 5723616 A	03-03-1998
US 6046333	A	04-04-2000	AU 4993497 A	22-05-1998
			WO 9818815 A1	07-05-1998
WO 9633189	A	24-10-1996	AU 5486396 A	07-11-1996
			EP 0823904 A1	18-02-1998
			WO 9633189 A1	24-10-1996
EP 1149583	A	31-10-2001	BR 0101456 A	04-12-2001
			CA 2344089 A1	13-10-2001
			EP 1149583 A2	31-10-2001
			US 2001041673 A1	15-11-2001